

An Introduction to Heisenberg Groups

Stephen Semmes

Actually, a number of related objects are often called “Heisenberg groups”, and they appear in several ways.

Let n be a positive integer, and let λ be a positive real number. For each x in $\mathbf{R}^n$, consider the linear operator T_x acting on the vector space of continuous complex-valued functions on $\mathbf{R}^n$ with compact support defined by

$$(1) \quad T_x(f)(w) = f(w - x).$$

In other words, $T_x(f)$ is the same as f translated by x , in the sense that $T_x(f)(w + x) = f(w)$. Thus T_0 is the identity operator, and

$$(2) \quad T_{x+x'} = T_x \circ T_{x'}$$

for all $x, x' \in \mathbf{R}^n$. Also, T_x is the identity operator only when $x = 0$, and thus the correspondence $x \mapsto T_x$ provides an isomorphic embedding of $\mathbf{R}^n$, as an abelian group, into the group of invertible linear operators on the vector space of continuous functions with compact support on $\mathbf{R}^n$ under the binary operation composition. Of course there are numerous other vector spaces of functions on $\mathbf{R}^n$ that one might use here.

For each y in $\mathbf{R}^n$ define a linear operator U_y on the same space of functions on $\mathbf{R}^n$ by

$$(3) \quad U_y(f)(w) = \exp(2\pi i \lambda y \cdot w) f(w),$$

where $y \cdot w$ denotes the usual inner product on $\mathbf{R}^n$, namely,

$$(4) \quad y \cdot w = \sum_{j=1}^n y_j w_j, \quad y = (y_1, \dots, y_n), \quad w = (w_1, \dots, w_n).$$

As before, U_y is equal to the identity operator if and only if $y = 0$, and

$$(5) \quad U_{y+y'} = U_y \circ U_{y'}$$

for all $y, y' \in \mathbf{R}^n$. Thus the correspondence $y \mapsto U_y$ also defines an isomorphic embedding of $\mathbf{R}^n$ as an abelian group into the group of invertible linear transformations on our vector space of functions. Note that the operator U_y can be described in terms of translations on the Fourier transform side, rather than directly, as for T_x .

How do these two families of operators interact? What is the subgroup of the group of invertible linear operators that they generate? If α is a complex number which satisfies $|\alpha| = 1$, let C_α denote the linear operator on our vector space of functions on $\mathbf{R}^n$ defined by

$$(6) \quad C_\alpha(f)(w) = \alpha f(w).$$

That is, we simply multiply the function f by the scalar α . The correspondence $\alpha \mapsto C_\alpha$ defines an isomorphic embedding of the abelian group of complex numbers with modulus 1, under multiplication, into the group of invertible linear operators on our vector space of functions. Clearly C_α commutes with T_x and U_y for all α, x , and y in their respective domains, but in general T_x and U_y do not commute. However, for all x, y in $\mathbf{R}^n$, $U_y \circ T_x$ can be written as $T_x \circ U_y \circ C_\alpha$ for a complex number α of modulus 1. Indeed, for any function f on $\mathbf{R}^n$,

$$(7) \quad \begin{aligned} T_x(U_y(f))(w) &= U_y(f)(w - x) \\ &= \exp(2\pi i \lambda y \cdot (w - x)) f(w - x) \\ &= \exp(-2\pi i \lambda y \cdot x) \exp(2\pi i \lambda y \cdot w) T_x(f)(w) \\ &= \exp(-2\pi i \lambda y \cdot x) U_y(T_x(f))(w), \end{aligned}$$

so that

$$(8) \quad U_y \circ T_x = T_x \circ U_y \circ C_\alpha \quad \text{with } \alpha = \exp(2\pi i \lambda y \cdot x).$$

It is not hard to check that the collection of linear operators on our vector space of functions on $\mathbf{R}^n$ of the form $T_x \circ U_y \circ C_\alpha$, where x, y range through all elements of $\mathbf{R}^n$ and α ranges through all complex numbers of modulus 1, forms a subgroup of the group of invertible linear operators on this vector space. This subgroup is the same as the subgroup generated by the T_x 's and U_y 's for $x, y \in \mathbf{R}^n$.

Here is a closely connected point. For this it is convenient to work with functions f which are continuously differentiable. If ν is an element of $\mathbf{R}^n$, then we write D_ν for the operator of directional differentiation in the direction

ν , i.e.,

$$(9) \quad D_\nu(f)(w) = \left. \frac{d}{dt} \right|_{t=0} f(w + t\nu).$$

If μ is a linear function on $\mathbf{R}^n$, so that $\mu(x) = x \cdot u$ for some u in $\mathbf{R}^n$ and all x in $\mathbf{R}^n$, then let us write M_μ for the operator of multiplication by μ ,

$$(10) \quad M_\mu(f)(w) = \mu(w) f(w).$$

In general D_ν and M_μ do not commute, and in fact

$$(11) \quad D_\nu(M_\mu f) - M_\mu(D_\nu f) = \mu(\nu) f$$

for all continuously differentiable functions f on $\mathbf{R}^n$, where the scalar $\mu(\nu)$ is the same as $D_\nu(\mu)$, which is a constant since μ is linear.

Let us define $H_n(\mathbf{R})$ as follows. First, as a set, $H_n(\mathbf{R})$ is equal to $\mathbf{R}^n \times \mathbf{R}^n \times \mathbf{R}$, i.e., the set of ordered triples (x, y, t) where x, y lie in $\mathbf{R}^n$ and t lies in $\mathbf{R}$. Next, we define a binary operation on $H_n(\mathbf{R})$ by

$$(12) \quad (x, y, t) \cdot (x', y', t') = (x + x', y + y', t + t' + x' \cdot y).$$

With respect to this operation, one can verify that $H_n(\mathbf{R})$ becomes a group, with $(0, 0, 0)$ as the identity element, and $(-x, -y, -t)$ the inverse of (x, y, t) .

Consider the correspondence

$$(13) \quad (x, y, t) \mapsto T_x \circ U_y \circ C_\alpha, \quad \alpha = \exp(2\pi i \lambda t).$$

This defines a mapping from $H_n(\mathbf{R})$ into invertible linear mappings acting on functions on $\mathbf{R}^n$, and one can verify that this is a group homomorphism. A triple (x, y, t) corresponds to the identity operator if and only if $x = y = 0$ and t is an integer multiple of $1/\lambda$.

We can define $H_n(\mathbf{Z})$ to be the set of triples (x, y, t) where x, y lie in $\mathbf{Z}^n$ and t lies in $\mathbf{Z}$. It is easy to see that this defines a subgroup of $H_n(\mathbf{R})$.

In addition to being a group, $H_n(\mathbf{R})$ is a smooth manifold of dimension $2n+1$, and the group operation is smooth. The quotient $H_n(\mathbf{R})/H_n(\mathbf{Z})$ makes sense not only as a set of cosets, but also as a compact smooth manifold without boundary of dimension $2n+1$. When $n=1$, this manifold has dimension 3, and is one of the basic building blocks discussed in [61].

Define a_j in $H_n(\mathbf{Z})$ for $j=1, \dots, n$ to be the triple (x, y, t) such that all components of x are equal to 0 except for the j th component, which is equal to 1, and such that $y=0$ and $t=0$. Similarly, define b_j in $H_n(\mathbf{Z})$ for

$j = 1, \dots, n$ to be the triple (x, y, t) such that $x = 0$, all components of y are equal to 0 except for the j th component, which is equal to 1, and $t = 0$. Define c in $H_n(\mathbf{Z})$ to be the triple $(0, 0, 1)$. It is easy to see that

$$(14) \quad a_1, \dots, a_n, b_1, \dots, b_n, c$$

generate $H_n(\mathbf{Z})$. Indeed, if k, l lie in $\mathbf{Z}^n$ and m lies in $\mathbf{Z}$, then (k, l, m) is the same as

$$(15) \quad a_1^{k_1} \dots a_n^{k_n} b_1^{l_1} \dots b_n^{l_n} c^m$$

in the group, where of course this expression is interpreted using the group operation.

For $i, j = 1, \dots, n$ we have that

$$(16) \quad a_i a_j = a_j a_i, \quad b_i b_j = b_j b_i$$

and

$$(17) \quad a_i c = c a_i, \quad b_j c = c b_j.$$

If $1 \leq i, j \leq n$ and $i \neq j$, then

$$(18) \quad a_i b_j = b_j a_i.$$

When $i = j$ we have in place of this

$$(19) \quad b_i a_i = a_i b_i c$$

for $i = 1, \dots, n$. These relations describe the group $H_n(\mathbf{Z})$ completely; every element of $H_n(\mathbf{Z})$ can be represented in a unique way as (15), and these relations are adequate to define the group operation in terms of this representation.

In addition to the group structure on $H_n(\mathbf{R})$, there is a natural family of *dilations*. Namely, for each positive real number r , define a mapping δ_r from $H_n(\mathbf{R})$ to itself by

$$(20) \quad \delta_r(x, y, t) = (r x, r y, r^2 t).$$

This is a one-to-one mapping of $H_n(\mathbf{R})$ onto itself, $\delta_r \circ \delta_s = \delta_{rs}$ for all $r, s > 0$, and δ_1 is the identity mapping on $H_n(\mathbf{R})$. It is easy to check that these dilations preserve the group structure on $H_n(\mathbf{R})$.

In connection with complex variables it is convenient to employ a slightly different formulation. Define $\widetilde{H}_n$ as a set as $\mathbf{C}^n \times \mathbf{R}$. The group operation on $\widetilde{H}_n$ is defined by

$$(21) \quad (z, t) \cdot (z', t') = \left(z + z', t + t' + 2 \operatorname{Im} \sum_{j=1}^n z_j \overline{z'_j} \right),$$

where $z = (z_1, \dots, z_n)$, $z' = (z'_1, \dots, z'_n)$, and $\bar{a}$ denotes the complex conjugate of a complex number a . As before, $(0, 0)$ is the identity element of $\widetilde{H}_n$, and $(-z, -t)$ is the inverse of (z, t) .

Put

$$(22) \quad U = \{(w, \sigma) \in \mathbf{C}^n \times \mathbf{C} : \operatorname{Im} \sigma > |w|^2\},$$

where $|w|^2 = \sum_{j=1}^n |w_j|^2$, as usual. Thus

$$(23) \quad \partial U = \{(w, \sigma) \in \mathbf{C}^n \times \mathbf{C} : \operatorname{Im} \sigma = |w|^2\}.$$

For each (z, t) in $\widetilde{H}_n$, define the mapping $A_{(z,t)}$ from $\overline{U}$ to itself by

$$(24) \quad A_{(z,t)}(w, \sigma) = \left(w + z, \sigma + t + i|z|^2 + 2i \sum_{j=1}^n w_j \overline{z_j} \right).$$

Because

$$(25) \quad |w + z|^2 = |w|^2 + 2 \operatorname{Re} \sum_{j=1}^n w_j \overline{z_j} + |z|^2,$$

one can check that $A_{(z,t)}(w, \sigma) \in U$ when $(w, \sigma) \in U$ and $A_{(z,t)}(w, \sigma) \in \partial U$ when $(w, \sigma) \in \partial U$. Also,

$$\begin{aligned} (26) \quad & A_{(z,t)}(A_{(z',t')}(w, \sigma)) \\ &= \left(w + z' + z, \sigma + t' + t + i|z'|^2 + i|z|^2 + 2i \sum_{j=1}^n w_j \overline{z'_j} + 2i \sum_{j=1}^n (w_j + z'_j) \overline{z_j} \right) \\ &= \left(w + z' + z, \sigma + t' + t + i|z'|^2 + i|z|^2 + 2i \sum_{j=1}^n w_j \overline{(z'_j + z_j)} + 2i \sum_{j=1}^n z'_j \overline{z_j} \right) \\ &= \left(w + z' + z, \sigma + t' + t + i|z' + z|^2 + 2i \sum_{j=1}^n w_j \overline{(z'_j + z_j)} - 2 \operatorname{Im} \sum_{j=1}^n z'_j \overline{z_j} \right) \\ &= \left(w + z + z', \sigma + t + t' + i|z + z'|^2 + 2i \sum_{j=1}^n w_j \overline{(z_j + z'_j)} + 2 \operatorname{Im} \sum_{j=1}^n z_j \overline{z'_j} \right) \\ &= A_{(z,t) \cdot (z',t')}(w, \sigma). \end{aligned}$$

In particular, $A_{(-z,-t)}$ is the inverse of $A_{(z,t)}$ as a mapping from $\overline{U}$ onto itself.

From the definition of $A_{(z,t)}$ it is clear that it is a *holomorphic* mapping of U to itself, and in fact a complex-affine mapping, since there are only w 's and σ 's in $A_{(z,t)}(w, \sigma)$, and none of their complex conjugates. Thus $A_{(z,t)}$ is in fact a *biholomorphic* mapping of U onto itself.

Let us abuse our notation a bit and use δ_r to denote the natural dilation by $r > 0$ on $\widetilde{H}_n$,

$$(27) \quad \delta_r(z, t) = (r z, r^2 t).$$

As before, δ_r is a one-to-one mapping of $\widetilde{H}_n$ onto itself, $\delta_r \circ \delta_s = \delta_{rs}$ for all $r, s > 0$, δ_1 is the identity mapping on $\widetilde{H}_n$, and δ_r preserves the group structure on $\widetilde{H}_n$. These dilations also correspond to mappings on $\overline{U}$ in a natural way. Namely, define $\Delta_r : \overline{U} \rightarrow \overline{U}$ for $r > 0$ by

$$(28) \quad \Delta_r(w, \sigma) = (r w, r^2 \sigma).$$

It is easy to see that Δ_r is a one-to-one mapping of $\overline{U}$ onto itself which takes U to U , and that $\Delta_r \circ \Delta_s = \Delta_{rs}$ for all $r, s > 0$ and Δ_1 is the identity mapping. Furthermore, Δ_r is a biholomorphic mapping of U onto itself, and

$$(29) \quad \Delta_r(A_{(z,t)}(w, \sigma)) = A_{\delta_r(z,t)}(\Delta_r(w, \sigma)).$$

A famous fact about U is that it is biholomorphically equivalent to the unit ball in $\mathbf{C}^{n+1}$. Thus the $A_{(z,t)}$'s and Δ_r 's correspond to biholomorphic transformations on the unit ball in $\mathbf{C}^{n+1}$. They do not account for all of the biholomorphic transformations, though; basically what is missing are the complex-linear transformations on $\mathbf{C}^{n+1}$ which map the unit ball onto itself, which is the same as saying that they preserve the standard Hermitian inner product on $\mathbf{C}^{n+1}$.

References

- [1] G. Alexopoulos, *Sub-Laplacians with Drift on Lie Groups of Polynomial Volume Growth*, *Memoirs of the American Mathematical Society* **739**, 2002.
- [2] L. Ambrosio and P. Tilli, *Selected Topics on "Analysis on Metric Spaces"*, Scuola Normale Superiore, Pisa, 2000.

- [3] R. Beals, B. Gaveau, and P. Greiner, *Hamilton–Jacobi theory and the heat kernel on Heisenberg groups*, Journal de Mathématiques Pures et Appliquées (9) **79** (2000), 633–689.
- [4] R. Beals and P. Greiner, *Calculus on Heisenberg Manifolds*, Annals of Mathematics Studies **119**, Princeton University Press, 1988.
- [5] R. Beals, P. Greiner, and N. Stanton, *The heat equation and geometry of CR manifolds*, Bulletin of the American Mathematical Society (N.S.) **10** (1984), 275–276.
- [6] R. Beals, P. Greiner, and N. Stanton, *The heat equation on a CR manifold*, Journal of Differential Geometry **20** (1984), 343–387.
- [7] A. Bellaïche and J.-J. Risler, editors, *Sub-Riemannian Geometry*, Birkhäuser, 1996.
- [8] C. Berenstein, D.-C. Chang, and J. Tie, *Laguerre Calculus and its Applications on the Heisenberg Group*, American Mathematical Society and International Press, 2001.
- [9] A. Borel, *Semisimple Groups and Riemannian Symmetric Spaces*, Hindustan Book Agency, 1998.
- [10] J. Burillo, *Lower bounds of isoperimetric functions for nilpotent groups*, in *Geometric and Computational Perspectives on Infinite Groups*, DIMACS Series in Discrete Mathematics and Theoretical Computer Science **25**, American Mathematical Society, 1996.
- [11] R. Coifman and G. Weiss, *Analyse Harmonique Non-Commutative sur Certains Espaces Homogènes*, Lecture Notes in Mathematics **242**, Springer-Verlag, 1971.
- [12] R. Coifman and G. Weiss, *Extensions of Hardy spaces and their use in analysis*, Bulletin of the American Mathematical Society **83** (1977), 569–645.
- [13] T. Coulhon and L. Saloff-Coste, *Isopérimétrie pour les groupes et les variétés*, Revista Matemática Iberoamericana **9** (1993), 293–314.
- [14] D. Danielli, N. Garofalo, and D.-M. Nhieu, *Notions of convexity in Carnot groups*, preprint, 2002.

- [15] D. Epstein, J. Cannon, D. Holt, S. Levy, M. Paterson, and W. Thurston, *Word Processing in Groups*, Jones and Bartlett, 1992.
- [16] B. Farb, *Automatic groups: A guided tour*, L'Enseignement Mathématique (2) **38** (1992), 291–313.
- [17] G. Folland, *Harmonic Analysis in Phase Space*, Annals of Mathematics Studies **122**, Princeton University Press, 1989.
- [18] G. Folland and E. Stein, *Hardy Spaces on Homogeneous Groups*, Mathematical Notes **28**, Princeton University Press, 1982.
- [19] B. Franchi, R. Serapioni, and F. Serra Cassano, *Sur les ensembles de périmètre fini dans le groupe de Heisenberg*, Comptes Rendus des Séances de l'Académie des Sciences Paris Série I Mathématiques **329** (1999), 183–188.
- [20] B. Franchi, R. Serapioni, and F. Serra Cassano, *Rectifiability and perimeter in the Heisenberg group*, Mathematische Annalen **321** (2001), 479–531.
- [21] B. Franchi, R. Serapioni, and F. Serra Cassano, *Regular hypersurfaces, intrinsic perimeter and implicit function theorem in Carnot groups*, to appear, Communications in Analysis and Geometry.
- [22] W. Goldman, *Complex Hyperbolic Geometry*, Oxford University Press, 1999.
- [23] P. Hajlasz and P. Koskela, *Sobolev Met Poincaré*, Memoirs of the American Mathematical Society **688**, 2000.
- [24] P. de la Harpe, *Topics in Geometric Group Theory*, University of Chicago Press, 2000.
- [25] J. Heinonen, *Calculus on Carnot groups*, in *Fall School in Analysis (Jyväskylä, 1994)*, 1–31, Reports of the Department of Mathematics, University of Jyväskylä **68**, 1995.
- [26] J. Heinonen, *Lectures on Analysis on Metric Spaces*, Springer-Verlag, 2001.

- [27] S. Helgason, *Differential Geometry, Lie Groups, and Symmetric Spaces*, Academic Press, 1978.
- [28] R. Howe, *On the role of the Heisenberg group in harmonic analysis*, Bulletin of the American Mathematical Society (N.S.) **3** (1980), 821–843.
- [29] M. Kapovich, *Hyperbolic Manifolds and Discrete Groups*, Birkhäuser, 2001.
- [30] S. Kobayashi, *Transformation Groups in Differential Geometry*, Springer-Verlag, 1972.
- [31] S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry, Volume II*, Wiley, 1969.
- [32] A. Koranyi, *The Poisson integral for generalized half-planes and bounded symmetric domains*, Annals of Mathematics (2) **82** (1965), 332–350.
- [33] A. Koranyi, *Harmonic functions on Hermitian hyperbolic space*, Transactions of the American Mathematical Society **135** (1969), 507–516.
- [34] A. Koranyi, *A remark on boundary values of functions of several variables*, in *Several Complex Variables I, Maryland 1970*, Lecture Notes in Mathematics **155**, Springer-Verlag, 1970.
- [35] A. Koranyi and H. Reimann, *Quasiconformal mappings on the Heisenberg group*, Inventiones Mathematicae **80** (1985), 309–338.
- [36] A. Koranyi and H. Reimann, *Foundation for the theory of quasiconformal mappings on the Heisenberg group*, Advances in Mathematics **111** (1995), 1–87.
- [37] A. Koranyi and S. Vagi, *Intégrales singulières sur certains espaces homogènes*, Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences Séries A et B **268** (1969), A765–A768.
- [38] A. Koranyi and S. Vagi, *Singular integrals in homogeneous spaces and some problems in classical analysis*, Annali della Scuola Normale Superiore di Pisa (3) **25** (1971), 575–648.

- [39] A. Koranyi and S. Vagi, *Cauchy–Szegő integrals for systems of harmonic functions*, Annali della Scuola Normale Superiore di Pisa (3) **26** (1972), 181–196.
- [40] A. Koranyi, S. Vagi, and G. Welland, *Remarks on the Cauchy integral and the conjugate function in generalized half-planes*, Journal of Mathematics and Mechanics **19** (1970) 1069–1081.
- [41] S. Krantz, *Function Theory of Several Complex Variables*, Wiley, 1982.
- [42] S. Krantz, *Complex Analysis: The Geometric Viewpoint*, Carus Mathematical Monographs **23**, Mathematical Association of America, 1990.
- [43] S. Krantz, *A Panorama of Harmonic Analysis*, Carus Mathematical Monographs **27**, Mathematical Association of America, 1999.
- [44] G.-P. Leonardi and S. Rigot, *Isoperimetric sets on Carnot groups*, to appear, Houston Journal of Mathematics.
- [45] G. Lu, J. Manfredi, and B. Stroffolini, *Convex functions on the Heisenberg group*, preprint, 2002.
- [46] V. Magnani, *Elements of Geometric Measure Theory on Sub-Riemannian Groups*, thesis, Scuola Normale Superiore, Pisa, 2002.
- [47] J. Mitchell, *On Carnot–Carathéodory metrics*, Journal of Differential Geometry **21** (1985), 35–45.
- [48] A. Nagel and E. Stein, *Lectures on Pseudodifferential Operators: Regularity Theorems and Applications to Nonelliptic Problems*, Mathematical Notes **24**, Princeton University Press, 1979.
- [49] P. Pansu, *Une inégalité isopérimétrique sur le groupe de Heisenberg*, Comptes Rendus des Séances de l’Académie des Sciences Série I Mathématiques **295** (1982), 127–130.
- [50] P. Pansu, *An isoperimetric inequality on the Heisenberg group*, Conference on Differential Geometry on Homogeneous spaces (Torino, 1983), Rendiconti del Seminario Matematico, Università e Politecnico di Torino, Special Issue (1983), 159–174.

- [51] P. Pansu, *Métriques de Carnot–Carathéodory et quasiisométries des espaces symétriques de rang un*, Annals of Mathematics (2) **129** (1989), 1–60.
- [52] W. Rudin, *Functional Analysis*, McGraw-Hill, 1973.
- [53] W. Rudin, *Function Theory in the Unit Ball of $\mathbf{C}^n$* , Springer-Verlag, 1980.
- [54] E. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, 1970.
- [55] E. Stein, *Boundary Behavior of Holomorphic Functions of Several Complex Variables*, Mathematical Notes **11**, Princeton University Press, 1972.
- [56] E. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton University Press, 1993.
- [57] E. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton University Press, 1971.
- [58] R. Strichartz, *Sub-Riemannian Geometry*, Journal of Differential Geometry **24** (1986), 221–263; corrections, **30** (1989), 595–596.
- [59] M. Taylor, *Noncommutative microlocal analysis I*, Memoirs of the American Mathematical Society **313**, 1984.
- [60] M. Taylor, *Noncommutative Harmonic Analysis*, Mathematical Surveys and Monographs **22**, American Mathematical Society, 1986.
- [61] W. Thurston, *Three-Dimensional Geometry and Topology*, edited by S. Levy, Princeton University Press, 1997.
- [62] S. Vagi, *Harmonic analysis on Cartan and Siegel domains*, in *Studies in Harmonic Analysis*, MAA Studies in Mathematics **13**, 257–309, Mathematical Association of America, 1976.
- [63] N. Varopoulos and S. Mustapha, *Analysis on Lie Groups*, Cambridge University Press, 2001.
- [64] N. Varopoulos, L. Saloff-Coste, and T. Coulhon, *Analysis and Geometry on Groups*, Cambridge University Press, 1992.